

**Press Release No. 20/2019**

Press Office  
Universitätsstrasse 10  
78464 Konstanz  
GERMANY  
+49 7531 88-3603  
Fax +49 7531 88-3766

kum@uni-konstanz.de  
www.uni-konstanz.de/en

19.03.2019

**New technique for in-cell distance determination**

**Researchers from the University of Konstanz, Bielefeld University and ETH Zurich demonstrate for the first time that the pulsed EPR technique RIDME (relaxation-induced dipolar modulation enhancement) can be used for in-cell distance determination in biomacromolecules. Applied within the cell, RIDME improves significantly on conventional double electron–electron resonance (DEER) measurements.**

In a joint paper which has just been published in *The Journal of Physical Chemistry Letters*, researchers from the University of Konstanz, Bielefeld University and ETH Zurich demonstrate for the first time that the electron paramagnetic resonance (EPR) technique RIDME (relaxation-induced dipolar modulation enhancement) can be applied to determine distances between gadolinium(III)-based spin labels in cells. In-cell distance determination by electron paramagnetic resonance (EPR) reveals essential structural information about biomacromolecules, including their conformation as well as folding and unfolding processes.

Conventional methods for in-cell determination of distances such as double electron–electron resonance (DEER or PELDOR) are principally much less sensitive than RIDME, provide up to five times smaller modulation depths, have certain limitations with regard to excitation bandwidth and are technically more demanding. As a single-frequency technique which makes use of relaxation-induced spin flips to determine the distance between two spin labels, i.e. between two unpaired electrons, RIDME overcomes all of these disadvantages.

What is special about this technique is that it allows the researchers to work with molecules under native conditions, as Professor Malte Drescher and lead author Dr Mykhailo Azarkh, both from the University of Konstanz, emphasize: “We started out by analysing the conformation of a protein inside the cell. With less sensitive techniques, we are forced to insert and tag a lot of protein to be able to observe it, which is not at all what happens in nature. Ideally, we want to be working with concentrations that are physiologically relevant. Since RIDME is much more sensitive than DEER, it allows us to do just that. We are now in a position to address issues that we would not otherwise be able to address”.

The performance of in-cell RIDME was assessed at Q-band using stiff molecular rulers labelled with Gd(III)-PyMTA and microjointed into *Xenopus laevis* (African clawed frog) oocytes. In other words, the researchers used a model system where the precise distance between the spin labels was already known, allowing them to verify the RIDME measurements. The resulting paper entitled “Gd(III)–Gd(III) Relaxation-Induced Dipolar Modulation Enhancement for In-Cell Electron Paramagnetic Resonance Distance Determination” was published online in *The Journal of Physical Chemistry Letters* on 13.03.2019.

In-cell RIDME distance determination was developed and tested as part of the on-going ERC-funded project “SPICE – Spectroscopy in cells”, for which Malte Drescher, Heisenberg Professor for Spectroscopy of Complex Systems at the University of Konstanz, and his research team were

recognized with an ERC Consolidator Grant worth approximately two million euros in 2017. Their goal is to develop new approaches to spectroscopy that allow them to explore larger and more complex biological structures at the molecular level of the cell.

The next step in this line of research will be to identify other suitable spin labels and to develop RIDME for application in molecules where the distance between the spin labels is unknown. A particular focus of attention will be on molecules associated with neuro-degenerative diseases such as Alzheimer's and Parkinson's.

**Facts:**

- Original publication: Mykhailo Azarkh, Anna Bieber, Mian Qi, Jörg W. A. Fischer, Maxim Yulikov, Adelheid Godt, Malte Drescher. Gd(III)–Gd(III) Relaxation-Induced Dipolar Modulation Enhancement for In-Cell Electron Paramagnetic Resonance Distance Determination. *J. Phys. Chem. Lett.* 2019, 10, pp 1477–1481. DOI: <https://doi.org/10.1021/acs.jpcllett.9b00340>.
- Researchers from the University of Konstanz, Bielefeld University and ETH Zurich demonstrate for the first time that the EPR technique RIDME (relaxation-induced dipolar modulation enhancement) can be used for in-cell determination of distances in biomacromolecules.
- As opposed to other techniques, such as double electron–electron resonance (DEER), RIDME is technically less demanding, has no limitations with respect to the excitation bandwidth and provides five times larger modulation depth. There are no artefacts caused by pseudosecular terms.
- The results were generated as part of the on-going ERC project SPICE (“Spectroscopy in cells”), for which Professor Malte Drescher, Heisenberg Professor for Spectroscopy of Complex Systems at the University of Konstanz, received an ERC Consolidator Grant 2017 in the amount of approximately two million euros for a period of five years (2018-2022).

**Note to editors:**

An image is available for download here: [https://cms.uni-konstanz.de/fileadmin/pi/filesserver/2019/Bilder/New\\_technique\\_for\\_in-cell\\_distance\\_determination.jpg](https://cms.uni-konstanz.de/fileadmin/pi/filesserver/2019/Bilder/New_technique_for_in-cell_distance_determination.jpg)

**Caption:** In-cell distance determination by EPR reveals essential structural information about biomacromolecules under native conditions. For the first time, the pulsed EPR technique RIDME (relaxation induced dipolar modulation enhancement) was utilized for distance measurements inside cells. It provides a five-times improved sensitivity as compared to the previously used double electron-electron resonance approach.

**Image credit:** Research group Professor Malte Drescher, University of Konstanz

**Contact:**

University of Konstanz  
Communications and Marketing  
Phone: + 49 7531 88-3603  
Email: [kum@uni-konstanz.de](mailto:kum@uni-konstanz.de)

- [uni.kn/en](http://uni.kn/en)